

P3.280: Early Clinical Outcomes of VRN101099 in ERBB2-Mutated Advanced Non-Small Cell Lung Cancer: a Case Series from a Phase 1a Study

S.Y. Lee¹, C.G. Kim², J.H. Kim¹, R. Cosman³, S.Y. Rha², J.H. Youk⁴, S-A. Im⁴, D. Yu⁵, S.Y. Zhang⁶

¹Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul/KR, ²Yonsei Cancer Center, Yonsei University College of Medicine, Seoul/KR, ³The Kinghorn Cancer Centre, St. Vincent's Hospital, Sydney, Garvan Institute of Medical Research, University of New South Wales, Sydney/AU, ⁴Seoul National University Hospital, Seoul National University College of Medicine, Cancer Research Institute, Seoul National University, Seoul/KR, ⁵Voronoi, Incheon/KR, ⁶Voronoi USA, Inc., Boston/MA/USA

BD: Jay Ahn, ahn@voronoi.io | CMO: Shuang Yin Zhang, y.zhang@voronoi.io

INTRODUCTION

Backgrounds

- HER2-activating mutations occur in ~2–3% of solid tumors¹ and are associated with poor prognosis and frequent brain metastases in NSCLC².
- Recently approved HER2 TKIs, zongertinib and sevabertinib, have demonstrated clinical activity in HER2-mutant NSCLC, particularly against kinase domain mutations³.
- VRN101099** is a covalent, CNS-penetrant, HER2 selective TKI with a distinct **degradation mechanism (P3.240)** active across HER2 domains and tumor types.
- Here we report early clinical activity across HER2 mutation domains and tumor types in the first HER2-mutant patients from the Phase 1a study (**NCT06806982**).

Methods

- Ten HER2-mutant patients had ≥1 post-baseline tumor assessment as of 2 September 2026. Patients A-D (most mature follow-up) and Patient E (an additional NSCLC case) are presented here.
- Patients received dose ranging from 80 to 480 mg with tumor assessment every 2 cycles according to RECIST v1.1 and for safety per CTCAE v5.0.





FIGURE 1. Map of Identified HER2 Mutations

CLINICAL CASES

Patient A 56 Female · 160 mg · HER2 S310Y · Lung Cancer


pembrolizumab+pemetrexed+cisplatin SD → **trastuzumab+neratinib** SD → **T-DXd** SD → **VRN101099 PR**



- Prior therapies, including T-DXd, all reached **only stable disease**
- Achieved PR at C5D1 with continued response by **-49.1% at C14D1**
- Overall PR** with **sequential CR** in **TLs; Lung at C3D1** and **Breast at C14D1**

Patient B 60 Male · 320 mg · HER2 L755P · Lung Cancer

MEDI5752(anti-PD-1xCTLA-4 bispecific antibody)+pemetrexed+carboplatin SD → pembrolizumab+pemetrexed+carboplatin PD → **T-DXd PR** → **VRN101099 PR**



- PR maintained** with **VRN101099** following prior **T-DXd (17 months)**
- Overall PR** with **-35.8%** tumor reduction in 2 Lung TL at C5D1
- G3 diarrhea** resolved within a week via anti-diarrheal regimen

Patient Characteristics

	Patient A	Patient B	Patient C	Patient D	Patient E
Age/Sex	56/F	60/M	46/F	72/M	62/F
ECOG	0	1	0	0	0
Dose (mg)	160	320	160	80	240
HER2 mutation	S310Y	L755P	V777L	S310F	A775_G776insYVMA
HER2 domain	ECD	TKD	TKD	ECD	TKD
Primary site	Lung	Lung	Breast	Pancreas	Lung
Metastatic site	Lymph Nodes, Brain , LT.PLEURAL	Lymph Nodes, Bone, Brain , Pleural seeding	Liver, Lung, Pleura	Gastric, Lymph node	Lung, Lymph Nodes, Brain
Prior HER2 Tx	trastuzumab, neratinib, T-DXd	T-DXd	T-DXd	trastuzumab, tucatinib	sevabertinib, ELVN-002, T-DXd
Prior Systemic Tx	3	3	7	3	7

ECD, Extracellular Domain; TKD, Tyrosine Kinase Domain

- Enrolled patients carried HER2 mutations in the **extracellular domain (ECD; S310Y, S310F)** and **kinase domain (L755P, V777L, A775_G776insYVMA)** in **lung, breast, and pancreas**.
- Heavily pretreated (median of 3 prior systemic therapies) with prior exposure to HER2-directed therapies: **4/5 post T-DXd, 3/5 post-HER2 TKI (neratinib, tucatinib)**.
- 3/5 had baseline brain metastases**.

Patient C 46 Female · 160 mg · HER2 V777L · Breast Cancer

palbociclib+letrozole **PR** → fulvestrant Unknown → fulvestrant+alpelisib **PD** → everolimus+exemestane SD → capecitabine **PR** → paclitaxel **PR** → **T-DXd PR** → **VRN101099 SD**

- Anti-Tumor activity in **heavily pre-treated** setting (**≥7 prior Tx**)
- Tumor response by RECIST v1.1: **SD (-29.7% in Liver TL at C5D1)**
- Discontinued due to **non-TL PD**


Patient D 72 Male · 80 mg · HER2 S310F · Pancreatic Cancer

FOLFIRINOX Unknown → nab-paclitaxel+gemcitabine Unknown → **trastuzumab+tucatinib** SD → **VRN101099 SD**

- Tumor response by RECIST v1.1: **SD (-26.7% shrinkage in Stomach at C3D1)**
- Tx interrupted for ~3 weeks due to underlying disease
- Discontinued to **subject withdrawal**

Patient E 62 Female · 240 mg · HER2 A775_G776insYVMA · Lung Cancer

pemetrexed+cisplatin **PD** → atezolizumab **PD** → sevabertinib **PR** → ELVN-002(HER2 TKI) SD → gemcitabine SD → docetaxel **PD** → **T-DXd PR** → **VRN101099 PR**



- 2 Lung TL PR** achieved at **first tumor assessment** of **-56.1%**
- Deep partial response** following relapse on **sevabertinib, ELVN-002, and T-DXd**
- Brain NTL** also decreased

CLINICAL & MOLECULAR OUTCOME

Depth of Response




FIGURE 2: Anti-Tumor Activity of VRN101099 in Patients with Diverse HER2-Mutated Cancers

Anti-tumor activity of VRN101099 extended beyond the kinase domain to include extracellular domain mutations and diverse histology. Among 10 response-evaluable HER2-mutant patients, ORR was 40.0% and DCR was 90.0%.

CDC, Gallbladder carcinoma; UC, Urothelial Carcinoma; CRC, Colorectal Cancer; PC, Pancreatic Cancer; BC, Breast Cancer; NSCLC, Non-Small Cell Lung Cancer; RCC, Renal Cell Carcinoma




FIGURE 3: Drug Exposure and Tumor Response

Exploratory analysis included baseline plasma ctDNA profiling (Guardant360) and serial pharmacokinetic sampling during VRN101099 treatment. Tumor shrinkage was correlated with higher drug exposure levels.

Genetic Profiling and Durability




FIGURE 4: Genomic Context, Response, and Treatment Duration

Durable PRs occurred in NSCLC patients with baseline brain metastases carrying both extracellular (S310Y) and kinase domain (L755P, A775_G776insYVMA) mutations regardless of co-occurring genetic alterations and high variant allele fraction .

SUMMARY

EFFICACY

Of 10 patients with HER2 activating mutations, 3 NSCLC patients with brain metastases showed efficacy. Despite heavy treatment history including T-DXd exposure just prior to enrollment, all three achieved PR with decreased intracranial non-target lesions.

COVERAGE

Antitumor activity extends across the mutation and tumor spectrum. VRN101099 exposure translated into tumor reduction from NSCLC to breast (TKD mutation) and pancreatic (ECD mutation) cancers.

NEXT STEPS

A Phase 1b expansion is planned with cohorts stratified by HER2 status (HER2-mutant NSCLC and HER2-positive solid tumors), including combinations with other HER2-directed agents.